

Emergent Kinetics and Fractionalized Charge in 1D Spin-Orbit Coupled Flatband Optical Lattices

Fei Lin,¹ Chuanwei Zhang,² and V. W. Scarola¹

¹*Department of Physics, Virginia Tech, Blacksburg, Virginia 24061, USA*

²*Department of Physics, The University of Texas at Dallas, Richardson, Texas 75080, USA*

(Received 29 August 2013; revised manuscript received 15 November 2013; published 18 March 2014)

Recent ultracold atomic gas experiments implementing synthetic spin-orbit coupling allow access to flatbands that emphasize interactions. We model spin-orbit coupled fermions in a one-dimensional flatband optical lattice. We introduce an effective Luttinger-liquid theory to show that interactions generate collective excitations with emergent kinetics and fractionalized charge, analogous to properties found in the two-dimensional fractional quantum Hall regime. Observation of these excitations would provide an important platform for exploring exotic quantum states derived solely from interactions.

DOI: 10.1103/PhysRevLett.112.110404

PACS numbers: 67.85.-d, 03.65.Vf, 03.75.Ss, 05.30.Fk

Introduction.—Emergent quantum states derived from interactions can exhibit rich structure because they are, by definition, not adiabatically connected to the underlying single-particle states. Two-dimensional (2D) electron gases placed in a strong magnetic field offer seminal examples. In the absence of a magnetic field, 2D electrons typically demonstrate Fermi-liquid behavior, but a strong magnetic field, the fractional quantum Hall (FQH) limit [1], would seem to prevent the formation of a Fermi liquid. This regime is defined by an absence of single-particle kinetic energy that leaves interparticle interactions to generate many-body quantum states in a flatband (the lowest Landau level). However, it is now well known that interesting properties, such as fractional charge from screening and other kinetic effects [2–4], emerge from interactions in the FQH regime. The remarkable fact that application of an external field first suppresses single-particle properties to leave interactions to generate similar emergent properties leads to a natural question: Can these emergent mechanisms manifest in other contexts? Flatbands in one dimension offer a logical analogue [5–8].

The Luttinger-liquid paradigm [9–12] captures the physics of many one-dimensional (1D) models. It predicts excitations with, e.g., fractionalized charge arising from competition between interactions and kinetic energy. External fields could, in analogy to 2D magnetic fields, be constructed to quench kinetics in one dimension, but the absence of kinetics in 1D flatbands would appear to rule out Luttinger-liquid behavior.

In this Letter, we show that kinetics, fractionalized charge excitations, and other Luttinger-liquid-like properties emerge solely from interactions in experimentally feasible 1D flatband models. Our proposal relies on recent experimental progress [13–18] in engineering synthetic spin-orbit coupling (SOC) for ultracold atomic gases [19]. These experiments show that Raman beams can be used to dress atoms with spin-dependent momentums. Rashba

(and/or Dresselhaus) SOC governing these dressed states [20,21] are tunable to extremes not possible in solids. Recent work shows that Rashba coupling in a 1D optical lattice [22] or gas [23,24] can be tuned to yield flatbands, a new limit that could play a role analogous to the lowest Landau level [25], but interaction effects in a 1D flat Rashba SOC band remain unexplored.

We study the impact of interactions between two-component fermions in a flat SOC band in 1D optical lattices. We find that the SOC elongates single-particle basis states to generate highly nontrivial nearest neighbor (NN) interactions [26]. The extended interactions lead to Wigner crystals of spinors with dispersive collective modes. These excitations are unexpected because they imply kinetics that emerge purely from interactions.

We predict that these excitations also exhibit fractionalized charge even in the flatband limit. To show this, we must contend with the fact that the absence of single-particle kinetic energy prevents direct application of the Luttinger-liquid theory. We find, instead, that emergent kinetics allows us to introduce an effective Luttinger-liquid theory. We compute the emergent velocities and fractionalized charge of excitations as experimentally verifiable observables. We also estimate the experimental parameters for observing these excitations. Detection of kinetics and fractionalized charge derived solely from interactions in one dimension would have important consequences for the study of emergent Luttinger-liquid behavior, in analogy to emergent fractional charge found in the 2D FQH regime.

Model.—We consider an equal population of N two-component fermions in a 1D optical lattice. We start with a noninteracting Hamiltonian that adds Rashba SOC to the optical lattice potential [22]:

$$\mathcal{H}_{\text{SOC}} = \frac{p_x^2}{2m} - sE_R \cos^2(k_L x) + \left(\frac{\hbar k_R}{m}\right) p_x \sigma_z + \Omega \sigma_x, \quad (1)$$

where p_x is the momentum of particles of mass m , s is the optical lattice strength, k_L is the optical lattice wave vector, $E_R = \hbar^2 k_L^2 / 2m$ is the recoil energy, $\hbar k_R / m$ is the SOC strength, σ are Pauli matrices, and Ω is the Zeeman field strength. We work in units of the lattice spacing, π / k_L .

Figure 1 plots the eigenvalues of \mathcal{H}_{SOC} , $\omega(k)$ to show that Eq. (1) yields flatbands. We project into the lowest flatband. Projection, achieved by considering only χ particles, is warranted in the presence of an energy gap between the χ and ζ bands at low densities [27].

We can derive a low-energy Hubbard model of interactions operating in such a flat Rashba band in the tight binding limit. In the Wannier basis, the interatom interactions (e.g., s -wave contact interactions between alkali atoms) become purely on site. After projection to the lowest flatband, the on-site Hubbard interaction defines the Hamiltonian of the entire system and is therefore the focus of our study [27]:

$$H = U \sum_{\{k\}} f_{\{k\}} \chi_{k_1}^\dagger \chi_{k_2}^\dagger \chi_{k_3} \chi_{k_4}, \quad (2)$$

where U is the on-site Hubbard repulsion that defines the only energy scale, χ_k^\dagger creates a fermion at wave vector k in the lowest band, and $f_{\{k\}} \equiv L^{-1} \delta_{k_1+k_2, k_3+k_4} \sin(\alpha_{k_1}) \times \cos(\alpha_{k_2}) \cos(\alpha_{k_3}) \sin(\alpha_{k_4})$. Here the Kronecker delta implies momentum conservation up to a reciprocal lattice vector, and L is the number of lattice sites.

Equation (2) is written in terms of lowest flatband-projected particles using a unitary transformation between the original fermions and flatband fermions so that




FIG. 1 (color online). (Main panels) Single-particle energy $\omega(k)$ of several lowest Bloch bands due to SOC for $s = 4$ (left) and $s = 10$ (right). The ratio of the lowest energy gap to the bandwidth is tuned to ≈ 8 [27] in both panels by setting $k_R = k_L/2$, $\Omega = 0.22E_R$ for $s = 4$, and $\Omega = 0.05E_R$ for $s = 10$. (Inset) Diagonal interaction between χ particles as a function of intersite distance, d , for $s = 4E_R$ and $s = 10E_R$ yielding $V_1/U \approx 0.0529$.

χ particles are defined as spinors of the original atoms [27]. We define the unitary transformation in terms of optical lattice parameters: $\tan(\alpha_k) = [\omega_1(k) - h(k)] / \Omega$ with $h(k) \equiv -2t \cos(k + k_R)$ and $\omega_1(k) \equiv -2t \cos k \cos k_R - \sqrt{4t^2 \sin^2 k \sin^2 k_R + \Omega^2}$. Here t is the NN hopping [27].

Projection to χ particles generates nontrivial delocalized single-particle basis states. To see this, we Fourier transform χ_k to real space. The on-site interaction between the original atoms becomes a longer range interaction between χ particles. The leading diagonal interaction, $V_d \tilde{n}_i \tilde{n}_{i+d}$, is between NN. Here $\tilde{n}_i \equiv \chi_i^\dagger \chi_i$.

The inset of Fig. 1 shows the interaction strength, V_d , between χ particles. Different optical lattice depths lead to the same interaction, where V_d falls off quickly with the dominant interaction given by V_1 , provided that the band remains flat [27]. The inset shows two key results: (1) The interaction is longer range, and (2) the form of the interaction is robust over a wide range of s . In the following, we can therefore focus on $s = 10$ without loss of generality.

Equation (2) contains a large number of terms, but by considering a few of the largest terms (with strengths V_1 , t_1^* , and t_2^*), we argue for intriguing low-energy states. Leading off-diagonal terms in Eq. (2) are given by conditional next nearest neighbor (NNN) hoppings of χ particles, i.e., $-|t_1^*| \chi_{i+2}^\dagger \tilde{n}_{i+1} \chi_i$ and $|t_2^*| \tilde{n}_i \chi_{i+3}^\dagger \chi_{i+1}$, where $|t_1^*|/U = 0.0257$ and $|t_2^*|/U = 0.0015$. We note that conditional hopping originates entirely from interactions. The V_1 term is the strongest and should generate crystal states of spinor χ particles, but conditional hoppings can give rise to emergent kinetics in excitations. We verify this picture below by combining diagonalization with an effective model.

Numerical results.—To more rigorously study Eq. (2), we use numerics to explore the low-energy Hilbert space and confine our study to half filling, $N/L = 1/2$. We note that the absence of kinetic energy excludes the direct use of Luttinger-liquid theory.

We numerically diagonalize Eq. (2) using the Lanczos algorithm. Translational symmetry allows us to work within a fixed total momentum sector, K . The left panel of Fig. 2 shows the four lowest total energies, $E(K)$, as a function of the total momentum for several system sizes. We find data collapse for $L \geq 16$. Our numerics therefore apply to the thermodynamic limit.

The ground state of H is a spinor Wigner crystal shown schematically in the left panel of Fig. 2, set to $E(\pm\pi/2) = 0$. Two Wigner crystals (both with particles at every other site) are defined in momentum space by a linear combination of wave functions at $K = \pm\pi/2$. We verify the crystalline nature of the ground state by breaking the degeneracy with a small, staggered chemical potential, $\mu \sum_i (-1)^i \tilde{n}_i$, added to Eq. (2). In the $\mu \rightarrow 0 \pm$ limit, the density shows that the system spontaneously picks one of the two degenerate Wigner crystal ground states [27]. We have also calculated the charge structure factor




FIG. 2 (color online). (Left panel) The many-body energy dispersion $E(K)$ (scattered symbols) for the flatband-projected Hamiltonian H on various lattice sizes, L . The solid curve is the fit of $E(K)$ with the effective extended Hubbard model [Eq. (4)] that highlights a dispersive collective mode. (Right panel) The energy dispersion of a similar classical model [Eq. (3)] with $V_1/U = 0.0529$, showing no dispersive collective modes. The four schematics denote representative ground and excited state configurations of spinor χ particles (encircled arrows) in real space.

$S(k) = L^{-2} \sum_{i,j} e^{ik(r_i - r_j)} \langle \tilde{n}_i \tilde{n}_j \rangle$. We find that $S(k)$ has well-defined peaks at $k = \pi$, indicating Wigner crystals.

We, for comparison, numerically solve a diagonal (classical) Hamiltonian known [28] to yield Wigner crystals:

$$H_D = V_1 \sum_i \tilde{n}_i \tilde{n}_{i+1}. \quad (3)$$

The right panel of Fig. 2 shows the many-body energy spectrum. The ground states of H_D coincide with those of Eq. (2), i.e., at $K = \pm\pi/2$, further showing that the ground states of Eq. (2) are classical Wigner crystals of spinor χ particles. The first excited state of H_D , however, is non-dispersive and lies at an energy V_1 . This is the energy cost of moving one particle in the Wigner crystal to a NN site (see the schematic of this classical excitation in Fig. 2, right panel). A comparison of the left and right panels shows that while the ground states are essentially the same, the excited states of Eq. (2) are fundamentally different from those of Eq. (3).

The excited states of Eq. (2), the left panel of Fig. 2, exhibit a gap $\sim 0.016U$ above the ground state. The conditional hopping terms cause the otherwise degenerate excited band to form a dispersive collective mode. The off-diagonal conditional hopping terms superpose the classical configurations of χ particles (see the schematic, left panel of Fig. 2). To better understand the nature of the excited states, we construct an effective model.

Effective Luttinger-liquid theory.—We construct an effective model of Eq. (2) by adding hopping terms to Eq. (3). The effective hopping terms are emergent because

they represent kinetics not present in the original model [Eq. (2)]. We verify the accuracy of the effective model by comparing energetics and by taking wave-function overlaps. The effective model is then studied using Luttinger-liquid theory on the emergent degrees of freedom.

We capture the effects of conditional hopping with ordinary single-particle hopping terms in an effective extended Hubbard model:

$$H_{\text{eff}} = - \sum_i [t_1 + t_2(-1)^i] (\chi_i^\dagger \chi_{i+2} + \text{H.c.}) + H_D, \quad (4)$$

where t_1 and t_2 are fitting parameters quantifying emergent NNN hopping. Figure 3 illustrates t_1 and t_2 in real space. Note that t_1 and t_2 scale with U because we added these parameters to capture the properties of excited states generated entirely by interactions in the original hopping-free model [Eq. (2)].

We vary t_1 and t_2 and numerically solve Eq. (4) to get the best fit of $E(K)$ while maximizing overlap of the corresponding wave functions. Table I shows representative ($L = 20$) fits for the lowest eigenstates. The energy differences between H_{eff} and H are all within 5%, and the wave-function overlaps for the lowest states are all above 50% with almost 100% overlaps at $K = 0$ and $\pm\pi$. The overlap between the ground states ($K = \pm\pi/2$) is above 99.9%. We plot the first excited state of Eq. (4) from the best fit parameters as the black curve in the left panel of Fig. 2 for comparison. The overlap and energetic comparison show that Eq. (4) captures the essential properties of Eq. (2) at low energies. We can therefore use Eq. (4) as an effective theory to make predictions for experiments.

We now show that Eq. (4) exhibits excitations with fractionalized charge quantified by Luttinger-liquid theory. We first diagonalize the hopping terms in Eq. (4) [27].




FIG. 3 (color online). (Main panel) Emergent single-particle energy dispersion [Eq. (5)]. The dashed and dotted lines cross at Fermi points, $k_F = \pm\pi/4$. Linearization is shown as solid straight lines. Differing slopes indicate asymmetric bands, i.e., $v_{1F} \neq v_{2F}$. (Inset) Schematic of hopping terms used in Eq. (4).

TABLE I. Fitting parameters (t_1 and t_2), the resulting energy differences ($\Delta E = E_{H_{\text{eff}}} - E_H$), and the wave-function overlaps between H_{eff} and H for an $L = 20$ system for the ground state at each total momentum sector, K , with $\mu/U = 10^{-5}$. Here the small energy differences and high wave-function overlaps indicate the quality of the effective model in capturing the essential physics of the original model.

$KL/2\pi$	t_1/U	t_2/U	$ \Delta E/E_H $	$\langle \Psi_{H_{\text{eff}}} \Psi_H \rangle$
0	0.0117	0.0130	0.035	0.99
1	0.0119	0.0135	0.010	0.97
2	0.0119	0.0135	0.010	0.88
3	0.0119	0.0135	0.015	0.74
4	0.0119	0.0135	0.005	0.53
5	0.0119	0.0135	0	0.99

The emergent “single-particle” energy dispersion has two energy bands ($b = 1, 2$):

$$\varepsilon_b(k) = -2[t_1 - (-1)^b t_2] \cos(2k), \quad (5)$$

with Fermi velocities $v_{bF} \equiv |\partial \varepsilon_b / \partial k|_{k_F} = |4[t_1 - (-1)^b t_2] \sin(2k_F)|$. For $N/L = 1/2$, each dispersion crosses the Fermi level at two Fermi points $k_F = \pm\pi/4$ (Fig. 3). Low-energy excitations near the Fermi points therefore consist of two left movers and two right movers.

We bosonize Eq. (4) to study interaction effects. We linearize the dispersion at the Fermi points [9,11,12], as depicted in Fig. 3. The elementary excitations near $\pm k_F$ are bosonic and, in the absence of the interacting term in Eq. (4), have the charge of the original flatband particles. We include H_D and find the normal modes of the bosonized Hamiltonian using a unitary transformation and rescaling of the bosonic fields [29]. The emergent normal mode Luttinger parameters, i.e., velocity, u_l , and the charge fractionalization ratio, g_l , are given by [27]

$$u_l = (v_{1F} v_{2F} \tilde{\lambda}_l)^{(1/2)}, \quad (6)$$

$$g_l = (\tilde{\lambda}_l)^{-(1/2)}, \quad (7)$$

where $l = 1, 2$ denotes the two normal modes, $\lambda_1 \equiv v_{1F}/v_{2F}$, $\lambda_2 \equiv \lambda_1^{-1}$, and $\tilde{\lambda}_l \equiv [\lambda_1 + \lambda_2 - (-1)^l \sqrt{(\lambda_1 - \lambda_2)^2 + 4V_1^2/(\pi^2 v_{1F} v_{2F})}]/2$. For $V_1 > 0$, we have $g_1 < 1$ indicating that the charge has fractionalized for this normal mode. To see this, we write the effective charge, q^* , in terms of the original charge, q , as $q^* = gq$ where q^* can be inferred from particle number conductance [10]. g_1 found here can be continuously tuned below unity. This should be contrasted with fractionally charged excitations in the FQH regime, where the fractions are only rational [2–4].

The Luttinger-liquid analysis therefore shows that low-energy collective modes of Eq. (4) can be thought of as fractionalized quasiparticles moving along a spinor Wigner

crystal. This result, while known in standard Luttinger-liquid theories [9–12], is surprising here since the single-particle eigenstates of the physical atoms are inert (flatband) particles that derive emergent kinetics from interactions. The close connection between Eqs. (4) and (2) also indicates that these modes should be experimentally observable.

Experimental requirements and observables.—Low temperatures and low atomic losses are, in general, difficult requirements for proposals to engineer strongly correlated quantum states with atomic gases. Most proposals require maximizing U by tuning a Feshbach resonance to enter strongly correlated regimes. However, Feshbach resonances contribute to unwanted heating and losses [30], particularly in SOC atomic gases [18,31]. The flatband regime studied here circumvents the need for strong U (and therefore a Feshbach resonance) because the system is automatically strongly correlated in the absence of kinetic energy.

We can estimate realistic parameters to show that the flatband regime is attainable. ^{40}K is one of the best candidates for strong SOC with low losses [18,31]. For ^{40}K in a 1D optical lattice with $s = 10$, $k_R = k_L/2$, $\Omega = 0.05E_R$, and a perpendicular confinement of lattice depth $60E_R$, we find $\Delta_{\text{SO}} \approx 0.10E_R$, $V_1 \approx 0.014E_R$ and $W \approx 0.013E_R$, where $W = 4t$ is the bandwidth. This shows that even bare s -wave scattering implies a strongly interacting flatband problem with $\Delta_{\text{SO}} \gg V_1 \gtrsim W$. Note that the last inequality is a very stringent flatband requirement. An accurate (but weaker) requirement assumes the many-body energy gap $\approx V_1/3$ (left panel, Fig. 2) to be larger than the single-particle hopping $V_1/3 \gtrsim W/4$. This implies that partial filling of the lowest band allows us to treat Eq. (1) as an irrelevant constant for realistic system parameters.

Parabolic confinement will compete with the many-body energy gap to diminish the size of the Wigner crystal near the trap center. The central crystal will give way to edge states when the parabolic trapping potential energy reaches the gap, i.e., $V_1/3 \approx m\omega_{\text{tr}}^2 x_{\text{max}}^2/2$. The crystal will then be as large as $\approx 2\sqrt{2V_1/3m\omega_{\text{tr}}^2}$ sites. Trapping potentials therefore place a lower bound on the size of the energy gap (and therefore U). For ^{40}K we find that even the bare s -wave scattering length allows significant crystal sizes, $\sim 86 - 150$ sites for realistic trapping strengths, $\omega_{\text{tr}} = 40 - 70$ Hz. Larger interaction strengths will increase the size of the crystal.

Observations of the states proposed here are in principle possible with currently available methods. The spinor Wigner crystal state manifests as a peak in the static structure factor of the original fermions, observable with demonstrated probes: noise correlations [32] or atomic matter wave scattering [33]. Luttinger-liquid parameters have also been observed by interfering Bose-Einstein condensates [34]. Detecting fractionalized charge is more

challenging. In the current context, it could be measured by, e.g., detection of partial backscattering from an impurity [10,35], optical methods [36,37], or charge pumping [38,39].

Summary.—We predict a set of intriguing collective states of matter in experiments with atomic Fermi gases confined to 1D optical lattices and in the presence of synthetic SOC. We constructed and studied a model where the atomic interactions operate in a flatband. We found that the single-particle basis states are delocalized spinors. Our analysis predicts that flatband spinor particles have surprising properties generated by on-site interactions among the original atoms: NN interactions and effective NNN hopping. The many-body ground state was found to be a Wigner crystal of spinors. We find that an effective Luttinger-liquid theory parametrizes emergent kinetics and fractionalized charge [Eq. (7)] in the low-energy collective modes, in direct analogy to the mechanism of emergence found in the FQH regime.

We acknowledge helpful comments from I. Spielman and support from the ARO (W911NF-12-1-0335, W911NF-12-1-0334), AFOSR (FA9550-11-1-0313), and DARPA-YFA.

-
- [1] D. C. Tsui, H. L. Stormer, and A. C. Gossard, *Phys. Rev. Lett.* **48**, 1559 (1982).
- [2] R. B. Laughlin, *Phys. Rev. Lett.* **50**, 1395 (1983).
- [3] S. D. Sarma and A. Pinczuk, *Perspectives in Quantum Hall Effects* (John Wiley & Sons, Inc., New York, 1997); J. K. Jain, *Composite Fermions* (Cambridge University Press, Cambridge, 2007).
- [4] V. J. Goldman and B. Su, *Science* **267**, 1010 (1995); R. de-Picciotto, M. Reznikov, M. Heiblum, V. Umansky, G. Bunin, and D. Mahalu, *Nature (London)* **389**, 162 (1997); L. Saminadayar, D. C. Glattli, Y. Jin, and B. Etienne, *Phys. Rev. Lett.* **79**, 2526 (1997).
- [5] E. Westerberg and T. H. Hansson, *Phys. Rev. B* **47**, 16554 (1993).
- [6] A. Seidel and D. H. Lee, *Phys. Rev. Lett.* **97**, 056804 (2006).
- [7] S. Jansen, E. H. Lieb, and R. Seiler, *Phys. Status Solidi (b)* **245**, 439 (2008).
- [8] M. Nakamura, Z. Y. Wang, and E. J. Bergholtz, *Phys. Rev. Lett.* **109**, 016401 (2012).
- [9] F. D. M. Haldane, *J. Phys. C* **14**, 2585 (1981).
- [10] M. P. A. Fisher and L. I. Glazman, in *Mesoscopic Electron Transport*, edited by L. L. Sohn, L. P. Kowenhoven, and G. Schon, *Proceedings of the NATO Advanced Study Institute* (Kluwer Academic Publishers, Dordrecht, The Netherlands, 1997).
- [11] E. Miranda, *Braz. J. Phys.* **33**, 3 (2003).
- [12] T. Gimarchy, *Quantum Physics in One Dimension* (Oxford University Press, New York, 2003).
- [13] Y.-J. Lin, K. Jimenez-Garcia, and I. B. Spielman, *Nature (London)* **471**, 83 (2011).
- [14] P. Wang, Z.-Q. Yu, Z. Fu, J. Miao, L. Huang, S. Chai, H. Zhai, and J. Zhang, *Phys. Rev. Lett.* **109**, 095301 (2012).
- [15] L. W. Cheuk, A. T. Sommer, Z. Hadzibabic, T. Yefsah, W. S. Bakr, and M. W. Zwierlein, *Phys. Rev. Lett.* **109**, 095302 (2012).
- [16] J. Y. Zhang *et al.*, *Phys. Rev. Lett.* **109**, 115301 (2012).
- [17] Z. Fu, L. Huang, Z. Meng, P. Wang, X.-J. Liu, H. Pu, H. Hu, and J. Zhang, *Phys. Rev. A* **87**, 053619 (2013).
- [18] R. A. Williams, M. C. Beeler, L. J. LeBlanc, K. Jimenez-Garcia, and I. B. Spielman, *Phys. Rev. Lett.* **111**, 095301 (2013).
- [19] I. Bloch, J. Dalibard, and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
- [20] J. D. Sau, R. Sensarma, S. Powell, I. B. Spielman, and S. Das Sarma, *Phys. Rev. B* **83**, 140510(R) (2011).
- [21] V. Galitski and I. B. Spielman, *Nature (London)* **494**, 49 (2013).
- [22] Y. Zhang and C. Zhang, *Phys. Rev. A* **87**, 023611 (2013).
- [23] X. Zhou, Y. Li, Z. Cai, and C. Wu, *J. Phys. B* **46**, 134001 (2013).
- [24] B. Ramachandhran, H. Hu, and H. Pu, *Phys. Rev. A* **87**, 033627 (2013).
- [25] V. W. Scarola, [arXiv:1305.6852](https://arxiv.org/abs/1305.6852).
- [26] R. A. Williams, L. J. LeBlanc, K. Jimenez-Garcia, M. C. Beeler, A. R. Perry, W. D. Phillips, and I. B. Spielman, *Science* **335**, 314 (2012).
- [27] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.112.110404> for a derivation of the flatband projected model and Luttinger parameters, and other calculation details.
- [28] J. Hubbard, *Phys. Rev. B* **17**, 494 (1978).
- [29] N. Sedlmayr, P. Korell, and J. Sirker, *Phys. Rev. B* **88**, 195113 (2013).
- [30] J. Stenger, S. Inouye, M. R. Andrews, H.-J. Miesner, D. M. Stamper-Kurn, and W. Ketterle, *Phys. Rev. Lett.* **82**, 2422 (1999).
- [31] R. Wei and E. J. Mueller, *Phys. Rev. A* **87**, 042514 (2013).
- [32] E. Altman, E. Demler, and M. D. Lukin, *Phys. Rev. A* **70**, 013603 (2004); S. Fölling, F. Gerbier, A. Widera, O. Mandel, T. Gericke, and I. Bloch, *Nature (London)* **434**, 481 (2005); I. B. Spielman, W. D. Phillips, and J. V. Porto, *Phys. Rev. Lett.* **98**, 080404 (2007).
- [33] B. Gadway, D. Pertot, J. Reeves, and D. Schneble, *Nat. Phys.* **8**, 544 (2012).
- [34] S. Hofferberth, I. Lesanovsky, T. Schumm, A. Imambekov, V. Gritsev, E. Demler, and J. Schmiedmayer, *Nat. Phys.* **4**, 489 (2008).
- [35] A. J. Daley, P. Zoller, and B. Trauzettel, *Phys. Rev. Lett.* **100**, 110404 (2008).
- [36] J. Ruostekoski, G. V. Dunne, and J. Javanainen, *Phys. Rev. Lett.* **88**, 180401 (2002).
- [37] J. Javanainen and J. Ruostekoski, *Phys. Rev. Lett.* **91**, 150404 (2003).
- [38] Y. Qian, M. Gong, and C. Zhang, *Phys. Rev. A* **84**, 013608 (2011).
- [39] L. Wang, M. Troyer, and X. Dai, *Phys. Rev. Lett.* **111**, 026802 (2013).